arXiv:1204.6064v1 [math.AP] 26 Apr 2012

Eigenfunction expansions for a fundamental solution of Laplace's equation on R^3 in parabolic and elliptic cylinder coordinates

H S Cohl[1] **and H Volkmer**[2]

[1]Information Technology Laboratory, National Institute of Standards and Technology, Gaithersburg, MD, USA

[2]Department of Mathematical Sciences, University of Wisconsin–Milwaukee, P. O. Box 413, Milwaukee, WI 53201, U.S.A.

E-mail: `hcohl@nist.gov`

Abstract. A fundamental solution of Laplace's equation in three dimensions is expanded in harmonic functions that are separated in parabolic or elliptic cylinder coordinates. There are two expansions in each case which reduce to expansions of the Bessel functions $J_0(kr)$ or $K_0(kr)$, $r^2 = (x - x_0)^2 + (y - y_0)^2$, in parabolic and elliptic cylinder harmonics. Advantage is taken of the fact that $K_0(kr)$ is a fundamental solution and $J_0(kr)$ is the Riemann function of partial differential equations on the Euclidean plane.

PACS numbers: 02.30.Em, 02.30.Gp, 02.30.Hq, 02.30.Jr, 02.30.Mv, 02.40.Dr

AMS classification scheme numbers: 35A08, 35J05, 42C15, 33E10, 33C15

1. Introduction

A fundamental solution of Laplace's equation

$$\frac{\partial^2 U}{\partial x^2}+\frac{\partial^2 U}{\partial y^2}+\frac{\partial^2 U}{\partial z^2}=0 \tag{1}$$

is given by (apart from a multiplicative factor of 4π)

$$U(\mathbf{x},\mathbf{x}_0)=\frac{1}{\|\mathbf{x}-\mathbf{x}_0\|},\quad \text{where } \mathbf{x}=(x,y,z)\neq\mathbf{x}_0=(x_0,y_0,z_0), \tag{2}$$

and $\|\mathbf{x}-\mathbf{x}_0\|$ denotes the Euclidean distance between $\mathbf{x}$ and $\mathbf{x}_0$. In many applications it is required to expand a fundamental solution in the form of a series or an integral, in terms of solutions of (1) that are separated in suitable curvilinear coordinates. Examples of such applications include electrostatics, magnetostatics, quantum direct and exchange Coulomb interactions, Newtonian gravity, potential flow, and steady state heat transfer. Morse & Feshbach (1953) [16] (see also Hobson (1955) [10]; MacRobert (1947) [12]; Heine (1881) [9]) provide a list of such expansions for various coordinate systems but the formulas for several coordinate systems are missing. It is the goal of this paper to provide these expansions for parabolic and elliptic cylinder coordinates. Although these expansions are partially known from Buchholz (1953) [2] and Lebedev (1972) [11] for parabolic cylinder coordinates, and from Meixner & Schäfke (1954) [14] for elliptic cylinder coordinates, we found it desirable to investigate these expansions in a systematic fashion and provide direct proofs for them based on eigenfunction expansions.

There will be two expansions for both of these coordinate systems. The first expansion for a cylindrical coordinate system on $\mathbf{R}^3$ starts from the known formula in terms of the integral of Lipschitz (see Watson (1944) [22, section 13.2]; Cohl *et al.* (2000) [5, (8)])

$$\frac{1}{\|\mathbf{x}-\mathbf{x_0}\|}=\int_0^\infty J_0(k\sqrt{(x-x_0)^2+(y-y_0)^2})e^{-k|z-z_0|}\,dk. \tag{3}$$

The Bessel function J_ν can be defined by (see for instance (10.2.2) in Olver *et al.* (2010) [17])

$$J_\nu(z):=\left(\frac{z}{2}\right)^\nu\sum_{n=0}^\infty\frac{(-z^2/4)^n}{n!\Gamma(\nu+n+1)}, \tag{4}$$

Note that for $W_k:\mathbf{R}^2\times\mathbf{R}^2\to\mathbf{R}$

$$W_k(x,y,x_0,y_0):=J_0(kr), \tag{5}$$

where $r^2:=(x-x_0)^2+(y-y_0)^2$, solves the partial differential equation

$$\frac{\partial^2 U}{\partial x^2}+\frac{\partial^2 U}{\partial y^2}+k^2U=0. \tag{6}$$

Therefore, in a cylindrical coordinate system on $\mathbf{R}^3$ involving the Cartesian coordinate z, the first expansion (3) reduces to expanding W_k from (5) in terms of solutions of (6) that are separated in a curvilinear coordinate system on the plane.

The second expansion for a cylindrical coordinate system on $\mathbf{R}^3$ is based on the known formula given in terms of the Lipschitz-Hankel integral (see Watson (1944) [22, section 13.21]; Cohl *et al.* (2000) [5, (9)])

$$\frac{1}{\|\mathbf{x}-\mathbf{x_0}\|}=\frac{2}{\pi}\int_0^\infty K_0(k\sqrt{(x-x_0)^2+(y-y_0)^2})\cos k(z-z_0)\,dk, \tag{7}$$

where $K_\nu:(0,\infty)\to\mathbf{R}$ (cf. (10.32.9) in Olver *et al.* (2010) [17]), the modified Bessel function of the second kind (Macdonald's function), of order $\nu\in\mathbf{R}$, is defined by

$$K_\nu(z):=\int_0^\infty e^{-z\cosh t}\cosh(\nu t)\,dt. \tag{8}$$

Now $V_k:\mathbf{R}^2\times\mathbf{R}^2\setminus\{(\mathbf{x},\mathbf{x}):\mathbf{x}\in\mathbf{R}^2\}\to(0,\infty)$, defined by

$$V_k(x,y,x_0,y_0):=K_0(kr) \tag{9}$$

solves the partial differential equation

$$\frac{\partial^2U}{\partial x^2}+\frac{\partial^2U}{\partial y^2}-k^2U=0. \tag{10}$$

In a cylindrical coordinate system on $\mathbf{R}^3$ involving the Cartesian coordinate z, the second expansion (7) reduces to expanding V_k in terms of solutions of (10) that are separated in curvilinear coordinates on the plane.

The paper is organized as follows. In section 2 we derive the desired expansion of $K_0(kr)$ in parabolic cylinder coordinates. This expansion is given in terms of series over Hermite functions. In section 3 we obtain an integral representation for $J_0(kr)$ in terms of separated solutions of (10) in terms of (modified) parabolic cylinder functions. We show how these results are based on a general expansion theorem in terms of solutions of the differential equation

$$-u''-\frac{1}{4}\xi^2u=\lambda u. \tag{11}$$

In sections 4 and 5 we derive the fundamental solution expansions in elliptic cylinder coordinates for $J_0(kr)$ and $K_0(kr)$, respectively.

Throughout this paper we rely on the following definitions. The set of natural numbers is given by $\mathbf{N}:=\{1,2,3,\ldots\}$, the set $\mathbf{N}_0:=\{0,1,2,\ldots\}=\mathbf{N}\cup\{0\}$, and the set $\mathbf{Z}:=\{0,\pm1,\pm2,\ldots\}$. The set $\mathbf{R}$ represents the real numbers and the set $\mathbf{C}$ represents the complex numbers.

2. Expansion of $K_0(kr)$ for parabolic cylinder coordinates

Parabolic coordinates on the plane (ξ,η) (see for instance Chapter 10 in Lebedev (1972) [11]) are connected to Cartesian coordinates (x,y) by

$$x=\frac{1}{2}(\xi^2-\eta^2),\quad y=\xi\eta, \tag{12}$$

where $\xi \in \mathbf{R}$ and $\eta \in [0, \infty)$. To simplify notation we will first set $k = 1$ in (9). Then V_1 satisfies

$$\frac{\partial^2 U}{\partial x^2} + \frac{\partial^2 U}{\partial y^2} - U = 0 \quad \text{if } (x, y) \neq (x_0, y_0). \tag{13}$$

Let (ξ, η), (ξ_0, η_0) be parabolic coordinates on $\mathbf{R}^2$ for (x, y) and (x_0, y_0), respectively. Then V_1 transforms to

$$v(\xi, \eta, \xi_0, \eta_0) := K_0(r(\xi, \eta, \xi_0, \eta_0)), \quad (\xi, \eta) \neq \pm(\xi_0, \eta_0)$$

and $r(\xi, \eta, \xi_0, \eta_0)$ is defined by

$$r^2 = \frac{1}{4}\left[(\xi + \xi_0)^2 + (\eta + \eta_0)^2\right]\left[(\xi - \xi_0)^2 + (\eta - \eta_0)^2\right], \quad r > 0. \tag{14}$$

Here and in the following we allow all $\xi, \eta, \xi_0, \eta_0 \in \mathbf{R}$ such that $(\xi, \eta) \neq \pm(\xi_0, \eta_0)$. From (13) or by direct computation we obtain that v solves the equation

$$\frac{\partial^2 u}{\partial \xi^2} + \frac{\partial^2 u}{\partial \eta^2} - (\xi^2 + \eta^2)u = 0. \tag{15}$$

Separating variables $u(\xi, \eta) = u_1(\xi)u_2(\eta)$ in (15), we obtain the ordinary differential equations

$$u_1'' + (2n + 1 - \xi^2)u_1 = 0, \tag{16}$$

$$u_2'' - (2n + 1 + \eta^2)u_2 = 0, \tag{17}$$

where we will use only $n \in \mathbf{N}_0$. Equations (16), (17) have the general solutions

$$u_1(\xi) = c_1 e^{-\xi^2/2} H_n(\xi) + c_2 e^{\xi^2/2} H_{-n-1}(i\xi),$$

$$u_2(\eta) = c_3 e^{\eta^2/2} H_n(i\eta) + c_4 e^{-\eta^2/2} H_{-n-1}(\eta),$$

where $H_\nu : \mathbf{C} \to \mathbf{C}$ is the Hermite function which can be defined in terms of Kummer's function of the first kind M as (cf. (10.2.8) in Lebedev (1972) [11])

$$H_\nu(z) := \frac{2^\nu \sqrt{\pi}}{\Gamma\left(\frac{1-\nu}{2}\right)} M\left(-\frac{\nu}{2}, \frac{1}{2}, z^2\right) - \frac{2^{\nu+1}\sqrt{\pi}}{\Gamma\left(-\frac{\nu}{2}\right)} z M\left(\frac{1-\nu}{2}, \frac{3}{2}, z^2\right),$$

and

$$M(a, b, z) := \sum_{n=0}^{\infty} \frac{(a)_n z^n}{(b)_n n!} \tag{18}$$

(see for instance (13.2.2) in Olver *et al.* (2010) [17]). Note that the Kummer function of the first kind is entire in z and a, and is a meromorphic function of b. The Hermite function is an entire function of both z and ν. If $\nu = n \in \mathbf{N}_0$ then $H_\nu(z)$ reduces to the Hermite polynomial of degree n.

We will expand v as a function of ξ in an orthogonal series of functions $e^{-\xi^2/2} H_n(\xi)$, $n \in \mathbf{N}_0$, so that the coefficients

$$f_n(\eta, \xi_0, \eta_0) := \int_{-\infty}^{\infty} v(\xi, \eta, \xi_0, \eta_0) e^{-\xi^2/2} H_n(\xi)\, d\xi,$$

are to be evaluated. We do this based on the observation that the function $(\xi, \eta) \mapsto v(\xi, \eta, \xi_0, \eta_0)$ is a fundamental solution of equation (15). It has logarithmic singularities at the points $\pm(\xi_0, \eta_0)$. Arguing as in Volkmer (1984) [21, Theorem 1.11], we obtain the following integral representation for solutions of (15).

Theorem 2.1. *Let $u \in C^2(\mathbf{R}^2)$ be a solution of (15). Let $(\xi_0, \eta_0) \in \mathbf{R}^2$, and let C be a a closed rectifiable curve on $\mathbf{R}^2$ which does not pass through $\pm(\xi_0, \eta_0)$, and let $n^\pm$ be the winding number of C with respect to $\pm(\xi_0, \eta_0)$. Then we have*

$$2\pi \left[n^+ u(\xi_0, \eta_0) + n^- u(-\xi_0, -\eta_0)\right] = \int_C (u\partial_2 v - v\partial_2 u)\, d\xi + (v\partial_1 u - u\partial_1 v)\, d\eta,$$

where ∂_1, ∂_2 denote partial derivatives with respect to ξ, η, respectively.

We apply Theorem 2.1 to the solution

$$u(\xi, \eta) = e^{-\xi^2/2} H_n(\xi) e^{\eta^2/2} H_n(i\eta),$$

of (15), and for C we take the positively oriented boundary of the rectangle $|\xi| \le \xi_1$, $|\eta| \le \eta_1$, where $|\xi_0| < \xi_1$, $|\eta_0| < \eta_1$, so $n^+ = n^- = 1$. Then let $\xi_1 \to \infty$ and note that the integrals over the vertical sides converge to 0. The integrals over the horizontal sides of the rectangle give the same contribution because the integrand changes sign when (ξ, η) is replaced by $(-\xi, -\eta)$. Therefore, we obtain, for $\eta = \eta_1 > |\eta_0|$,

$$2\pi u(\xi_0, \eta_0) = \int_{-\infty}^{\infty} (v(\xi, \eta, \xi_0, \eta_0)\partial_2 u(\xi, \eta) - u(\xi, \eta)\partial_2 v(\xi, \eta, \xi_0, \eta_0)) d\xi. \tag{19}$$

Set $f(\eta) = f_n(\eta, \xi_0, \eta_0)$ and $g(\eta) = e^{\eta^2/2} H_n(i\eta)$. Then (19) can be written as

$$2\pi u(\xi_0, \eta_0) = g'(\eta) f(\eta) - g(\eta) f'(\eta). \tag{20}$$

By differentiating both sides of (20) with respect to η we see that f satisfies (17). Since $f(\eta)$ goes to 0 as $\eta \to \infty$, it follows that

$$f(\eta) = c e^{-\eta^2/2} H_{-n-1}(\eta),$$

where c is a constant. Going back to (20), we find that

$$2\pi u(\xi_0, \eta_0) = cW,$$

where $W = i^n$ is the (constant) Wronskian of $e^{-\eta^2/2} H_{-n-1}(\eta)$ and $g(\eta)$. Therefore, if $\eta > |\eta_0|$, we obtain

$$f_n(\eta, \xi_0, \eta_0) = 2\pi(-i)^n e^{-\xi_0^2/2} H_n(\xi_0) e^{\eta_0^2/2} H_n(i\eta_0) e^{-\eta^2/2} H_{-n-1}(\eta).$$

Using the well-known series expansion in terms of Hermite functions (Lebedev (1972) [11, Theorem 2, page 71]), we obtain the following result.

Theorem 2.2. *For $\xi, \eta, \xi_0, \eta_0 \in \mathbf{R}$ with $\eta > |\eta_0|$,*

$$K_0(r(\xi, \eta, \xi_0, \eta_0)) = \sqrt{\pi} e^{(\eta_0^2 - \xi_0^2 - \eta^2 - \xi^2)/2} \sum_{n=0}^{\infty} \frac{(-i)^n}{2^{n-1} n!} H_n(\xi) H_{-n-1}(\eta) H_n(\xi_0) H_n(i\eta_0),$$

where r is defined by (14).

The special case $\xi_0 = \eta_0 = 0$ of Theorem 2.2 can be found in [11, Problem 7, page 298]. If we multiply each ξ, η, ξ_0, η_0 by $\sqrt{k}$ then we obtain an expansion for $K_0(kr)$. Inserting this expansion into (7) yields our final result.

Theorem 2.3. *Let* $\mathbf{x}$, $\mathbf{x}_0$ *be points on* $\mathbf{R}^3$ *with parabolic cylinder coordinates* (ξ,η,z) *and* (ξ_0,η_0,z_0), *respectively. If* $\eta_{\lessgtr} := {\min \atop \max}\{\eta,\eta_0\}$ *then*

$$\frac{1}{\|\mathbf{x}-\mathbf{x}_0\|} = \frac{2}{\sqrt{\pi}}\int_0^\infty e^{-k/2(\xi^2+\eta^2+\xi_0^2-\eta_0^2)}\cos k(z-z_0)$$

$$\times\sum_{n=0}^\infty \frac{(-i)^n}{2^{n-1}n!}H_n(\sqrt{k}\xi)H_{-n-1}(\sqrt{k}\eta_>)H_n(\sqrt{k}\xi_0)H_n(i\sqrt{k}\eta_<)\,dk,$$

where $\mathbf{x}\neq\mathbf{x}_0$.

If we reverse the order of the infinite series and the definite integral in the above expression we obtain

$$\frac{1}{\|\mathbf{x}-\mathbf{x}_0\|} = \frac{2}{\sqrt{\pi}}\sum_{n=0}^\infty \frac{(-i)^n}{2^{n-1}n!}\int_0^\infty e^{-k/2(\xi^2+\eta^2+\xi_0^2-\eta_0^2)}$$

$$\times H_n(\sqrt{k}\xi)H_{-n-1}(\sqrt{k}\eta)H_n(\sqrt{k}\xi_0)H_n(i\sqrt{k}\eta_0)\cos k(z-z_0)\,dk.$$

It would be interesting to know the value of the above definite integral.

3. Expansion of $J_0(kr)$ for parabolic cylinder coordinates

Transforming equation (6) to parabolic coordinates (12) we obtain

$$\frac{\partial^2 u}{\partial\xi^2}+\frac{\partial^2 u}{\partial\eta^2}+k^2(\xi^2+\eta^2)u=0.$$

In order to simplify notation we will (temporarily) set $k=\frac{1}{2}$ and $\zeta=i\eta$ with ζ real. Thus we consider

$$\frac{\partial^2 u}{\partial\xi^2}-\frac{\partial^2 u}{\partial\zeta^2}+\frac{1}{4}(\xi^2-\zeta^2)u=0,\quad \xi,\zeta\in\mathbf{R}. \tag{21}$$

If $u_1(\xi)$ and $u_2(\zeta)$ are solutions of the ordinary differential equation (11) for some λ then $u(\xi,\zeta)=u_1(\xi)u_2(\zeta)$ solves (21).

The function $W_{1/2}$ from (5) transformed to (ξ,ζ) becomes

$$w(\xi,\zeta,\xi_0,\zeta_0)=J_0\left(\frac{1}{2}\tilde{r}(\xi,\zeta,\xi_0,\zeta_0)\right), \tag{22}$$

where $\tilde{r}^2$ is a symmetric polynomial defined by

$$\begin{aligned}4\tilde{r}^2 &:= \left[(\xi-\xi_0)^2-(\zeta-\zeta_0)^2\right]\left[(\xi+\xi_0)^2-(\zeta+\zeta_0)^2\right]\\ &= 8\xi\xi_0\zeta\zeta_0+\xi^4+\xi_0^4+\zeta^4+\zeta_0^4-2\xi_0^2\zeta^2-2\xi_0^2\zeta_0^2-2\zeta^2\zeta_0^2-2\xi^2\xi_0^2-2\xi^2\zeta^2-2\xi^2\zeta_0^2,\end{aligned}$$

and J_0 is the order zero Bessel function of the first kind (see (4)). For fixed $\zeta,\xi_0,\zeta_0\in\mathbf{R}$ consider the function $f:\mathbf{R}\to\mathbf{R}$ defined by

$$f(\xi):=w(\xi,\zeta,\xi_0,\zeta_0). \tag{23}$$

We wish to expand this function in terms of (modified) parabolic cylinder harmonics according to a general expansion theorem that is derived in the following subsection.

3.1. Spectral theory of (modified) parabolic cylinder harmonics – A singular Sturm-Liouville problem

We discuss the Sturm-Liouville problem

$$-u''-\tfrac{1}{4}x^2u=\lambda u,\quad -\infty<x<\infty, \tag{24}$$

involving the spectral parameter λ subject to

$$u\in L^2(-\infty,\infty).$$

By replacing $\frac{1}{4}x^2$ by $-\frac{1}{4}x^2$, we obtain the equation describing the harmonic oscillator whose eigenfunctions (Hermite functions) and the corresponding spectral theory is well-known. The spectral problem associated with (24) is far less known.

A discussion of differential equation (24) and its solutions can be found in [15] and in section 8.2 of (24) Erdélyi *et al.* (1982) [7] (see also Magnus (1941) [13]; Wells & Spence (1945) [23]; Cherry (1948) [3]; Darwin (1949) [6]).

We will follow Chapter 9 in Coddington & Levinson (1955) [4]. First note that, by [4, Corollary 2, page 231], equation (24) is in the limit-point case at $x=\pm\infty$. Therefore, one can apply section 5 of Chapter 9 in [4].

For $\lambda,x\in\mathbf{C}$ we define the functions $u_1(\lambda,x)$, $u_2(\lambda,x)$ as the solutions of (24) uniquely determined by the initial conditions

$$u_1(\lambda,0)=u_2'(\lambda,0)=1,\quad u_1'(\lambda,0)=u_2(\lambda,0)=0.$$

These functions may be expressed in terms of Kummer's function of the first kind (18)

$$u_1(\lambda,x)=e^{-\frac{i}{4}x^2}M\left(\tfrac{1}{4}+\tfrac{i}{2}\lambda,\tfrac{1}{2},\tfrac{i}{2}x^2\right), \tag{25}$$

$$u_2(\lambda,x)=e^{-\frac{i}{4}x^2}xM\left(\tfrac{3}{4}+\tfrac{i}{2}\lambda,\tfrac{3}{2},\tfrac{i}{2}x^2\right). \tag{26}$$

For $x>0$, the function

$$\begin{aligned}u_3(\lambda,x)&=e^{-\frac{i}{4}x^2}U\left(\tfrac{1}{4}+\tfrac{i}{2}\lambda,\tfrac{1}{2},\tfrac{i}{2}x^2\right) \\ &=\frac{\sqrt{\pi}}{\Gamma(\frac{3}{4}+\frac{i}{2}\lambda)}u_1(\lambda,x)-(1+i)\frac{\sqrt{\pi}}{\Gamma(\frac{1}{4}+\frac{i}{2}\lambda)}u_2(\lambda,x)\end{aligned} \tag{27}$$

is another solution of (24). Here the Kummer function of the second kind $U:\mathbf{C}\times\mathbf{C}\times(\mathbf{C}\setminus(-\infty,0])\to\mathbf{C}$ can be defined as (see Olver *et al.* (2010) [17, (13.2.42)])

$$U(a,b,z):=\frac{\Gamma(1-b)}{\Gamma(a-b+1)}M(a,b,z)+\frac{\Gamma(b-1)}{\Gamma(a)}z^{1-b}M(a-b+1,2-b,z).$$

Except when $z=0$, each branch of U is entire in a and b. We assume that $U(a,b,z)$ has its principal value. The asymptotic behavior of the Kummer function of the second kind [17, (13.2.6)] shows that $u_3(\lambda,\cdot)\in L^2(0,\infty)$ provided that $\Im\lambda<0$. Since (24) is in the limit-point case at $+\infty$, u_3 is the only solution with this property except for a constant factor. The Titchmarsh-Weyl functions $m_{\pm\infty}(\lambda)$, $\Im\lambda\neq 0$, are defined by the property that $u_1(\lambda,x)+m_{\pm\infty}(\lambda)u_2(\lambda,x)$ is square-integrable at $x=\pm\infty$. Therefore,

$$m_\infty(\lambda)=\begin{cases}-(1+i)\dfrac{\Gamma(\frac{3}{4}+\frac{i}{2}\lambda)}{\Gamma(\frac{1}{4}+\frac{i}{2}\lambda)} & \text{if } \Im\lambda<0,\\[2ex] -(1-i)\dfrac{\Gamma(\frac{3}{4}-\frac{i}{2}\lambda)}{\Gamma(\frac{1}{4}-\frac{i}{2}\lambda)} & \text{if } \Im\lambda>0.\end{cases}$$

By symmetry, we have

$$m_{-\infty}(\lambda) = -m_\infty(\lambda).$$

Using the notation of [4, Theorem 5.1, page 251], on obtains

$$M_{11}(\lambda) = \frac{-1}{2m_\infty(\lambda)}, \quad M_{12} = M_{21} = 0, \quad M_{22}(\lambda) = \frac{m_\infty(\lambda)}{2}.$$

Now using [4, page 250, last line], we find that, for $\lambda \in \mathbf{R}$,

$$\rho_1'(\lambda) := \rho_{11}'(\lambda) = \frac{1}{\pi} \lim_{\epsilon \to 0+} \Im\left(M_{11}(\lambda + i\epsilon)\right) = \frac{e^{\frac{\pi}{2}\lambda}}{4\sqrt{2}\pi^2} \left|\Gamma(\tfrac14 + \tfrac{i}{2}\lambda)\right|^2,$$

since [17, (5.4.5)]

$$\Gamma(\tfrac14 + iy)\Gamma(\tfrac34 - iy) = \frac{\pi\sqrt{2}}{\cosh(\pi y) + i\sinh(\pi y)}.$$

Moreover, $\rho_{12}(\lambda) = \rho_{21}(\lambda) = 0$ and

$$\rho_2'(\lambda) := \rho_{22}'(\lambda) = \frac{1}{\pi} \lim_{\epsilon \to 0+} \Im\left(M_{22}(\lambda + i\epsilon)\right) = \frac{e^{\frac{\pi}{2}\lambda}}{2\sqrt{2}\pi^2} \left|\Gamma(\tfrac34 + \tfrac{i}{2}\lambda)\right|^2.$$

Since the ρ-functions are real-analytic functions (with no jumps), we see that the spectrum of (24) is the whole real line $\mathbf{R}$ and there are no eigenvalues. The latter also follows from the known asymptotic behavior of the solutions of (24) (see [17, Chapter 12]).

Using a variant of Stirling's formula (see (5.11.9) in Olver *et al.* (2010) [17])

$$|\Gamma(x+iy)| \sim \sqrt{2\pi}|y|^{x-1/2}e^{-\pi|y|/2} \quad \text{as } x, y \in \mathbf{R},\ |y| \to \infty, \tag{28}$$

we can determine the asymptotic behavior of ρ_j', namely

$$\rho_1'(\lambda) \sim \frac{1}{2\pi}|\lambda|^{-1/2}e^{\pi(\lambda - |\lambda|)/2} \quad \text{as } |\lambda| \to \infty, \tag{29}$$

$$\rho_2'(\lambda) \sim \frac{1}{2\pi}|\lambda|^{1/2}e^{\pi(\lambda - |\lambda|)/2} \quad \text{as } |\lambda| \to \infty. \tag{30}$$

Applying [4, Theorem 5.2, page 251] to the analysis above, we obtain the following result on the spectral resolution associated with equation (24).

Theorem 3.1. *For a given function $f \in L^2(\mathbf{R})$, form the functions*

$$g_j(\lambda) = \int_{-\infty}^{\infty} u_j(\lambda, x) f(x)\, dx, \quad j = 1, 2,\ \lambda \in \mathbf{R}. \tag{31}$$

Then

$$g_j \in L^2(\mathbf{R}, \rho_j), \quad j = 1, 2,$$

or, equivalently,

$$\int_{-\infty}^{\infty} |g_j(\lambda)|^2 \rho_j'(\lambda)\, d\lambda < \infty, \quad j = 1, 2.$$

The function f can be represented in the form

$$f(x) = \sum_{j=1}^{2} \int_{-\infty}^{\infty} u_j(\lambda, x) g_j(\lambda) \rho_j'(\lambda)\, d\lambda. \tag{32}$$

Moreover, we have Parseval's equation

$$\int_{-\infty}^{\infty} |f(x)|^2\,dx = \sum_{j=1}^{2} \int_{-\infty}^{\infty} |g_j(\lambda)|^2 \rho_j'(\lambda)\,d\lambda.$$

Equations (31), (32) establish a one-to-one correspondence between f and (g_1, g_2). The integrals appearing in (31), (32) have to be interpreted in the L^2-sense. For instance, (31) means that

$$\int_{-n}^{n} u_j(\lambda, x) f(x) dx$$

converges to $g_j(\lambda)$ in $L^2(\mathbf{R}, \rho_j)$ as $n \to \infty$. Of course, if

$$\int_{-\infty}^{\infty} |u_j(\lambda, x) f(x)|\,dx < \infty \quad \text{for every } \lambda \in \mathbf{R},$$

then (31) is also true pointwise.

3.2. Applying the spectral theory to the expansion of $J_0(kr)$ in parabolic cylinder coordinates

Since $f(\xi) = O(|\xi|^{-1})$ in (23) as $|\xi| \to \infty$, we have that $f \in L^2(\mathbf{R})$. Therefore, we can expand f using Theorem 3.1, according to (32). For $\lambda \in \mathbf{R}$, we form the integrals

$$g_j(\lambda, \zeta, \xi_0, \zeta_0) = \int_{-\infty}^{\infty} w(\xi, \zeta, \xi_0, \zeta_0) u_j(\lambda, \xi)\,d\xi, \quad j = 1, 2. \tag{33}$$

These are absolutely convergent integrals because $f(\xi) = O(|\xi|^{-1})$ and $u_j(\lambda, \xi) = O(|\xi|^{-1/2})$ as $|\xi| \to \infty$.

Since $w(\xi, \zeta, \xi_0, \zeta_0)$ from (22) solves (21) for fixed (ξ_0, ζ_0), and $u_j(\lambda, \xi)$ from (25), (26) solves (24), it follows from differentiation under the integral sign followed by integration by parts (see for instance Schäfke (1963) [18, Satz 8, page 26]) that g_j solves (24) as a function of ζ. Since the function w is symmetric in its four variables, it is also true that g_j solves (24) as function of ξ_0 for fixed ζ_0, ζ and as function of ζ_0 for fixed ξ_0, ζ. From these properties of g_j it follows easily that there are functions $c_{jk\ell m} : \mathbf{R} \to \mathbf{R}$ with $j, k, \ell, m = 1, 2$, depending on λ but not on ζ, ξ_0, ζ_0 such that

$$g_j(\lambda, \zeta, \xi_0, \zeta_0) = \sum_{k,\ell,m=1}^{2} c_{jk\ell m}(\lambda) u_k(\lambda, \zeta) u_\ell(\lambda, \xi_0) u_m(\lambda, \zeta_0). \tag{34}$$

This formula holds for all $\lambda, \zeta, \xi_0, \zeta_0 \in \mathbf{R}$. Substituting $\zeta = \xi_0 = \zeta_0 = 0$ in (33), (34), we obtain

$$c_{j111}(\lambda) = \int_{-\infty}^{\infty} J_0(\tfrac{1}{4}\xi^2) u_j(\lambda, \xi)\,d\xi.$$

If $j = 2$, we integrate over an odd function, so $c_{2111}(\lambda) = 0$. By differentiating (34) with respect to ζ and/or ξ_0 and/or ζ_0 and then substituting $\zeta = \xi_0 = \zeta_0 = 0$ we find (after some calculations)

$$c_{jk\ell m}(\lambda) = \begin{cases} c_1(\lambda) & \text{if } j = k = \ell = m = 1, \\ c_2(\lambda) & \text{if } j = k = \ell = m = 2, \\ 0 & \text{otherwise.} \end{cases}$$

Here $c_j : \mathbf{R} \to \mathbf{R}$ for $j = 1, 2$ is given by (see Appendix A)

$$c_1(\lambda) := \int_{-\infty}^{\infty} J_0(\tfrac{1}{4}\xi^2) u_1(\lambda, \xi)\, d\xi = \frac{2\sqrt{2}\pi e^{-\pi\lambda/2}}{\cosh(\pi\lambda)|\Gamma\left(\frac{3}{4} + \frac{i\lambda}{2}\right)|^2}, \tag{35}$$

$$c_2(\lambda) := -\int_{-\infty}^{\infty} \xi^{-1} J_1(\tfrac{1}{4}\xi^2) u_2(\lambda, \xi)\, d\xi = \frac{-4\sqrt{2}\pi e^{-\pi\lambda/2}}{\cosh(\pi\lambda)|\Gamma\left(\frac{1}{4} + \frac{i\lambda}{2}\right)|^2}. \tag{36}$$

According to (32),

$$w(\xi, \zeta, \xi_0, \zeta_0) = \sum_{j=1}^{2} \int_{-\infty}^{\infty} c_j(\lambda) \rho_j'(\lambda) u_j(\lambda, \xi) u_j(\lambda, \zeta) u_j(\lambda, \xi_0) u_j(\lambda, \zeta_0) d\lambda. \tag{37}$$

Note that

$$c_1(\lambda)\rho_1'(\lambda) = \frac{1}{2\pi \cosh(\pi\lambda)} \left| \frac{\Gamma\left(\frac{1}{4} + i\frac{\lambda}{2}\right)}{\Gamma\left(\frac{3}{4} + i\frac{\lambda}{2}\right)} \right|^2 = \frac{1}{4\pi^3} \left|\Gamma\left(\tfrac{1}{4} + \tfrac{i\lambda}{2}\right)\right|^4, \tag{38}$$

$$c_2(\lambda)\rho_2'(\lambda) = -\frac{2}{\pi \cosh(\pi\lambda)} \left| \frac{\Gamma\left(\frac{3}{4} + i\frac{\lambda}{2}\right)}{\Gamma\left(\frac{1}{4} + i\frac{\lambda}{2}\right)} \right|^2 = \frac{-1}{\pi^3} \left|\Gamma\left(\tfrac{3}{4} + \tfrac{i\lambda}{2}\right)\right|^4, \tag{39}$$

where we used [17, (5.4.4), (5.5.5)]. It follows from (28) that

$$c_1(\lambda)\rho_1'(\lambda) \sim \frac{1}{\pi|\lambda| \cosh(\pi\lambda)} \quad \text{as } |\lambda| \to \infty, \tag{40}$$

$$c_2(\lambda)\rho_2'(\lambda) \sim -\frac{|\lambda|}{\pi \cosh(\pi\lambda)} \quad \text{as } |\lambda| \to \infty. \tag{41}$$

It is known (see Atkinson (1964), [1, (8.2.5)]) that, for fixed $x \in \mathbf{C}$, there is a constant C such that $u_j(\lambda, x) = O(e^{C|\lambda|^{1/2}})$. It follows from (40), (41), that the integrands in (37) decay exponentially and therefore the corresponding integrals are absolutely convergent. By the identity theorem for analytic functions we see that equation (37) is true for all $\xi, \zeta, \xi_0, \zeta_0 \in \mathbf{C}$.

After setting $\zeta = i\eta$ and $\zeta_0 = i\eta_0$ in (37), one obtains the following result.

Theorem 3.2. *Let $\xi, \eta, \xi_0, \eta_0 \in \mathbf{R}$. Then*

$$J_0\left(\frac{1}{2} r(\xi, \eta, \xi_0, \eta_0)\right) = \sum_{j=1}^{2} \int_{-\infty}^{\infty} c_j(\lambda)\rho_j'(\lambda) u_j(\lambda, \xi) u_j(\lambda, i\eta) u_j(\lambda, \xi_0) u_j(\lambda, i\eta_0)\, d\lambda,$$

where r is given by (14) and $c_j(\lambda)\rho_j'(\lambda)$ is given by (38), (39).

In the special case $\xi_0 = \eta_0 = 0$ (or correspondingly $\xi = \eta = 0$), Theorem 3.2 can be found in Buchholz (1953) [2, (16), page 175]. Of course, if we multiply each ξ, η, ξ_0, η_0 by $\sqrt{2k}$ we get the expansion of $J_0(k\, r(\xi, \eta, \xi_0, \eta_0))$. This leads to the $J_0(kr)$ expansion of a fundamental solution for the three-dimensional Laplace equation in parabolic cylindrical coordinates.

Theorem 3.3. *Let* $\mathbf{x}$*,* $\mathbf{x}_0$ *be points on* $\mathbf{R}^3$ *with parabolic coordinates* (ξ, η, z) *and* (ξ_0, η_0, z_0)*, respectively, Then*

$$\frac{1}{\|\mathbf{x}-\mathbf{x}_0\|} = \sum_{j=1}^{2} \int_0^\infty \int_{-\infty}^\infty c_j(\lambda)\rho_j'(\lambda)$$
$$\times\, u_j(\lambda, 2\sqrt{k}\xi)u_j(\lambda, 2i\sqrt{k}\eta)u_j(\lambda, 2\sqrt{k}\xi_0)u_j(\lambda, 2i\sqrt{k}\eta_0)e^{-k|z-z_0|}\, d\lambda\, dk.$$

3.3. The Riemann method of integration

The function $w(\xi, \zeta, \xi_0, \zeta_0)$ (as a function of (ξ, ζ)) is a solution of the partial differential equation (21) and it satisfies the condition $w(\xi, \zeta, \xi_0, \zeta_0) = 1$ if $\xi - \xi_0 = \pm(\zeta - \zeta_0)$. This shows that w is the Riemann function of (21); for instance, see Garabedian (1986) [8].

The Riemann method of integration applied to the partial differential equation (21) as in Volkmer (1980) [19] gives, for all $\zeta, \xi_0, \zeta_0 \in \mathbf{R}$,

$$2u(\xi_0, \zeta_0) = u(\zeta - \zeta_0 + \xi_0, \zeta) + u(-\zeta + \zeta_0 + \xi_0, \zeta) \tag{42}$$
$$\int_{-\zeta+\zeta_0+\xi_0}^{\zeta-\zeta_0+\xi_0} [w(\xi, \zeta, \xi_0, \zeta_0)\partial_2 u(\xi, \zeta) - \partial_2 w(\xi, \zeta, \xi_0, \zeta_0)u(\xi, \zeta)]\, d\xi,$$

where $u \in C^2(\mathbf{R}^2)$ is a solution of (21). This formula for $\xi_0 = \zeta = 0$ and $u(\xi, \zeta) = u_1(\lambda, \xi)u_2(\lambda, \zeta)$ with u_1, u_2 from (25), (26) (after replacing ζ_0 by ζ), implies that

$$\int_{-\zeta}^{\zeta} J_0(\tfrac{1}{4}(\xi^2 - \zeta^2))u_1(\lambda, \xi)\, d\xi = 2u_2(\lambda, \zeta). \tag{43}$$

This equation allows us to transform the even solution u_1 into the odd solution u_2 of equation (24). One can prove (43) directly by denoting the left-hand side of (43) by $f(\zeta)$ and then showing that f is an odd solution of (24) with $f'(0) = 2$.

By differentiating (42) first with respect to ξ_0, ζ_0, and using $w_1 = u_2$, $w_2 = u_1$, we obtain (after a lengthy calculation) that

$$\int_{-\zeta}^{\zeta} \frac{\xi\zeta}{\xi^2 - \zeta^2} J_1(\tfrac{1}{4}(\xi^2 - \zeta^2))u_2(\lambda, \xi)\, d\xi = 2u_1(\lambda, \zeta) - 2u_2'(\lambda, \zeta).$$

This formula can also be proved directly.

Let $h : \mathbf{R}^2 \to \mathbf{R}$ be the function defined by

$$h(\xi, \zeta) := \begin{cases} J_0\left(\frac{1}{4}(\xi^2 - \zeta^2)\right) & \text{if } |\xi| < |\zeta|, \\ 0 & \text{otherwise.} \end{cases}$$

For fixed ζ this is an even function in $L^2(\mathbf{R})$ which can be expanded according to Theorem 3.1 (without knowing $c_j(\lambda)$), so that

$$\operatorname{sign}(\zeta)h(\xi, \zeta) = 2\int_{-\infty}^{\infty} u_1(\lambda, \xi)u_2(\lambda, \zeta)\rho_1'(\lambda)\, d\lambda.$$

If $\xi = 0$, we obtain

$$\operatorname{sign}(\zeta)J_0(\tfrac{1}{4}\zeta^2) = 2\int_{-\infty}^{\infty} u_2(\lambda, \zeta)\rho_1'(\lambda)\, d\lambda.$$

By Theorem 3.1, this formulas allows us to conclude

$$\int_0^\infty J_0(\tfrac14\zeta^2)u_2(\lambda,\zeta)\,d\zeta=\frac{\rho_1'(\lambda)}{\rho_2'(\lambda)},$$

and this is in agreement with (A.5).

4. Expansion of $J_0(kr)$ for elliptic cylinder coordinates

Consider equation (6) for $k>0$ and elliptic coordinates on the plane

$$x=c\cosh\xi\cos\eta,\quad y=c\sinh\xi\sin\eta, \tag{44}$$

where $\xi\in[0,\infty)$, $\eta\in\mathbf{R}$, and $c>0$. Transforming $u(\xi,\eta)=U(x,y)$ we obtain

$$\frac{\partial^2u}{\partial\xi^2}+\frac{\partial^2u}{\partial\eta^2}+k^2c^2(\cosh^2\xi-\cos^2\eta)u=0. \tag{45}$$

Separating variables $u(\xi,\eta)=u_1(\xi)u_2(\eta)$, leads to

$$-u_1''(\xi)+(\lambda-2q\cosh 2\xi)u_1(\xi)=0, \tag{46}$$

$$u_2''(\eta)+(\lambda-2q\cos 2\eta)u_2(\eta)\ =0, \tag{47}$$

where $q=\frac14c^2k^2>0$.

For $q\in\mathbf{R}$, Mathieu's equation (47) is a Hill's differential equation with period π; see [17] or [14]. Here q is positive but in the next section q will be negative. As a Hill's equation, Mathieu's equation (47) admits nontrivial 2π-periodic solutions if and only if λ is equal to one of its eigenvalues $a_n(q)$, $n\in\mathbf{N}_0$ or $b_n(q)$, $n\in\mathbf{N}$. If $\lambda=a_n(q)$, then Mathieu's equation has an even 2π-periodic solution $\mathrm{ce}_n(\eta,q)$, and if $\lambda=b_n(q)$ then Mathieu's equation has an odd 2π-periodic solution $\mathrm{se}_n(\eta,q)$. These functions are normalized according to

$$\int_0^{2\pi}\mathrm{ce}_n^2(\eta,q)\,d\eta=\int_0^{2\pi}\mathrm{se}_n^2(\eta,q)\,d\eta=\pi.$$

Moreover,

$$\mathrm{ce}_n(\eta+\pi,q)=(-1)^n\mathrm{ce}(\eta,q),\quad \mathrm{se}_n(\eta+\pi,q)=(-1)^n\mathrm{se}(\eta,q).$$

Note that all solutions of Mathieu's equation are entire functions of η.

For these 2π-periodic solutions of the Mathieu equation, we have the following expansion theorem.

Theorem 4.1. *Let $f(z)$ be a 2π-periodic function that is analytic in an open doubly-infinite strip S that contains the real axis. Then*

$$f(z)=\alpha_0\mathrm{ce}_0(z,q)+\sum_{n=1}^\infty(\alpha_n\mathrm{ce}_n(z,q)+\beta_n\mathrm{se}_n(z,q)), \tag{48}$$

where

$$\alpha_n=\frac1\pi\int_{-\pi}^{\pi}f(x)\mathrm{ce}_n(x,q)\,dx,\quad \beta_n=\frac1\pi\int_{-\pi}^{\pi}f(x)\mathrm{se}_n(x,q)\,dx.$$

The series (48) converges absolutely and uniformly on any compact subset of the strip S.

Let (x_0, y_0) and (x, y) be points on $\mathbf{R}^2$ with distance r. Let (x_0, y_0), (x, y) have elliptic coordinates (ξ_0, η_0) and (ξ, η), respectively. Then

$$r^2 = c^2 \left[(\cosh \xi \cos \eta - \cosh \xi_0 \cos \eta_0)^2 + (\sinh \xi \sin \eta - \sinh \xi_0 \sin \eta_0)^2\right]. \tag{49}$$

Clearly, $J_0(kr)$ as a function of (ξ, η) solves (45). We substitute $\zeta = i\xi$ and $\zeta_0 = i\xi_0$. Then (45) changes to

$$\frac{\partial^2 u}{\partial \zeta^2} - \frac{\partial^2 u}{\partial \eta^2} + k^2 c^2 (\cos^2 \eta - \cos^2 \zeta) u = 0, \tag{50}$$

and $J_0(kr)$ is transformed to

$$w(\zeta, \eta, \zeta_0, \eta_0) = J_0(k\tilde{r}),$$

where

$$\begin{aligned} \tilde{r}^2 &= c^2[(\cos \zeta \cos \eta - \cos \zeta_0 \cos \eta_0)^2 - (\sin \zeta \sin \eta - \sin \zeta_0 \sin \eta_0)^2] \\ &= c^2(\cos(\zeta - \eta) - \cos(\zeta_0 - \eta_0))(\cos(\zeta + \eta) - \cos(\zeta_0 + \eta_0)). \end{aligned}$$

The function $w(\zeta, \eta, \zeta_0, \eta_0)$ is an analytic function on $\mathbf{C}^4$ and it solves equation (50) as a function of (ζ, η). It is the Riemann function of this differential equation because $w(\zeta, \eta, \zeta_0, \eta_0) = 1$ if $\zeta - \zeta_0 = \pm(\eta - \eta_0)$. For fixed η, ζ_0, η_0 we wish to expand the function $\zeta \mapsto w(\zeta, \eta, \zeta_0, \eta_0)$ in a series of Mathieu functions according to Theorem 4.1. To this end we have to evaluate the integral

$$\int_{-\pi}^{\pi} w(\zeta, \eta, \zeta_0, \eta_0) \mathrm{ce}_n(\zeta, q)\, d\zeta,$$

and a similar integral with ce_n replaced by se_n. Using Riemann's method of integration applied to a pentagonal curve, it has been shown in [20] that

$$\int_{-\pi}^{\pi} w(\zeta, \eta, \zeta_0, \eta_0) \mathrm{ce}_n(\zeta, q)\, d\zeta = \mu_n(q) \mathrm{ce}_n(\eta, q) \mathrm{ce}_n(\zeta_0, q) \mathrm{ce}_n(\eta_0, q), \tag{51}$$

$$\int_{-\pi}^{\pi} w(\zeta, \eta, \zeta_0, \eta_0) \mathrm{se}_n(\zeta, q)\, d\zeta = \nu_n(q) \mathrm{se}_n(\eta, q) \mathrm{se}_n(\zeta_0, q) \mathrm{se}_n(\eta_0, q). \tag{52}$$

There do not exist explicit formulas for the quantities $\mu_n(q)$ and $\nu_n(q)$ but they can be determined as follows. Mathieu's equation (47) with $\lambda = a_n(q)$, $n \in \mathbf{N}_0$, has the solution $u_1(\eta) = \mathrm{ce}_n(\eta, q)$. We choose a second linear independent solution $u_2(\eta)$. Then there is σ such that

$$u_2(\eta + \pi) = \sigma u_1(\eta) + (-1)^n u_2(\eta)$$

and

$$\mu_n(q) = \frac{2(-1)^n \sigma}{W[u_1, u_2]}, \tag{53}$$

where $W[u_1, u_2]$ denotes the Wronskian of u_1 and u_2. Similarly, Mathieu's equation (47) with $\lambda = b_n(q)$, $n \in \mathbf{N}$, has the solution $u_3(\eta) = \mathrm{se}_n(\eta, q)$. We choose a second linear independent solution $u_4(\eta)$. Then there is τ such that

$$u_4(\eta + \pi) = \tau u_3(\eta) + (-1)^n u_4(\eta)$$

and

$$\nu_n(q) = \frac{2(-1)^n\tau}{W[u_3,u_4]}. \tag{54}$$

Now applying Theorem 4.1 and substituting $\zeta = i\xi$, $\zeta_0 = i\xi_0$, we obtain the following result.

Theorem 4.2. *Let* $\xi, \eta, \xi_0, \eta_0 \in \mathbf{C}$, *and let* $k > 0$, $c > 0$, $q = \frac{1}{4}c^2k^2$. *Then*

$$\begin{aligned} J_0(kr) &= \frac{1}{\pi}\sum_{n=0}^{\infty} \mu_n(q)\mathrm{ce}_n(i\xi,q)\mathrm{ce}_n(\eta,q)\mathrm{ce}_n(i\xi_0,q)\mathrm{ce}_n(\eta_0,q) \\ &\quad + \frac{1}{\pi}\sum_{n=1}^{\infty} \nu_n(q)\mathrm{se}_n(i\xi,q)\mathrm{se}_n(\eta,q)\mathrm{se}_n(i\xi_0,q)\mathrm{se}_n(\eta_0,q), \end{aligned}$$

where r *is given by (49).*

Theorem 4.2 agrees with expansion (23) (for $j = 1$ and $\nu = 0$), section 2.66 in Meixner & Schäfke [14], who have a slightly different notation. They use $\mathrm{me}_n(z,q)$, $n \in \mathbf{Z}$, where

$$\begin{aligned} \mathrm{me}_n(z,q) &:= \sqrt{2}\mathrm{ce}_n(z,q) \quad \text{if } n \in \mathbf{N}_0, \\ \mathrm{me}_{-n}(z,q) &:= -\sqrt{2}i\mathrm{se}_n(z,q) \quad \text{if } n \in \mathbf{N}. \end{aligned}$$

Moreover, the coefficients $\mu_n(q)$ and $\nu_n(q)$ are represented in a different form. The proof of Theorem 4.2 based on Riemann's method of integration appears to be new.

We now use (3) to obtain our final result in this section.

Theorem 4.3. *Let* $\mathbf{x}$, $\mathbf{x}_0$ *be points on* $\mathbf{R}^3$ *with elliptic cylinder coordinates* (ξ,η,z) *and* (ξ_0,η_0,z_0), *respectively. Then*

$$\begin{aligned} \frac{1}{\|\mathbf{x}-\mathbf{x}_0\|} &= \frac{1}{\pi}\sum_{n=0}^{\infty}\int_0^\infty \mu_n(q)\mathrm{ce}_n(i\xi,q)\mathrm{ce}_n(\eta,q)\mathrm{ce}_n(i\xi_0,q)\mathrm{ce}_n(\eta_0,q)e^{-k|z-z_0|}\,dk \\ &\quad + \frac{1}{\pi}\sum_{n=1}^{\infty}\int_0^\infty \nu_n(q)\mathrm{se}_n(i\xi,q)\mathrm{se}_n(\eta,q)\mathrm{se}_n(i\xi_0,q)\mathrm{se}_n(\eta_0,q)e^{-k|z-z_0|}\,dk, \end{aligned}$$

where $q = \frac{1}{4}c^2k^2$.

5. Expansion of $K_0(kr)$ for elliptic cylinder coordinates

Consider equation (10) for $k > 0$. Transforming to elliptic coordinates (44), we obtain

$$\frac{\partial^2 u}{\partial \xi^2} + \frac{\partial^2 u}{\partial \eta^2} - k^2c^2(\cosh^2\xi - \cos^2\eta)u = 0. \tag{55}$$

Separating variables $u(\xi,\eta) = u_1(\xi)u_2(\eta)$, leads again to (46), (47) but now $q = -\frac{1}{4}c^2k^2$ is negative.

We will need the following solutions of the modified Mathieu equation (46) when $q < 0$; see [17, §28.20]. Set $q = -h^2$ with $h > 0$. For $n \in \mathbf{N}_0$, $\mathrm{Ie}_n(\xi,h)$ is the even solution of (46) with $\lambda = a_n(q)$ with asymptotic behavior

$$\mathrm{Ie}_n(\xi,h) \sim I_n(2h\cosh\xi) \quad \text{as } \xi \to +\infty,$$

while $\mathrm{Ke}_n(\xi,h)$ is the recessive solution determined by

$$\mathrm{Ke}_n(\xi,h)\sim K_n(2h\cosh\xi)\quad \text{as } \xi\to+\infty,$$

where $I_n(z):=i^{-n}J_n(iz)$ and $K_n(z)$ are the modified Bessel functions of the first [17, (10.27.6)] and second kinds respectively, with integer order n (see (4), (8)). Similarly, for $n\in\mathbf{N}$, $\mathrm{Io}_n(\xi,h)$ is the odd solution of (46) with $\lambda=b_n(q)$ with asymptotic behavior

$$\mathrm{Io}_n(\xi,h)\sim I_n(2h\cosh\xi)\quad \text{as } \xi\to+\infty,$$

while $\mathrm{Ko}_n(\xi,h)$ is the recessive solution determined by

$$\mathrm{Ko}_n(\xi,h)\sim K_n(2h\cosh\xi)\quad \text{as } \xi\to+\infty.$$

For fixed $\xi,\xi_0,\eta_0\in\mathbf{R}$ we wish to expand the function

$$v(\xi,\eta,\xi_0,\eta_0):=K_0(kr(\xi,\eta,\xi_0,\eta_0))$$

with r given by (49) into a series of periodic Mathieu functions according to Theorem 4.1. The corresponding integrals appearing in the expansion will be computed based on the observation that $(\xi,\eta)\mapsto v(\xi,\eta,\xi_0,\eta_0)$ is a fundamental solution of (55). In fact, it is a solution of (55) and it has logarithmic singularities at the points $\pm(\xi_0,\eta_0+2m\pi)$, where m is any integer. Arguing as in Volkmer (1984) [21, Theorem 1.11], we have the following representation theorem for a solution of (55).

Theorem 5.1. *Let $u\in C^2(\mathbf{R}^2)$ be a solution of (55). Let $(\xi_0,\eta_0)\in\mathbf{R}^2$, and let C be a a closed rectifiable curve on $\mathbf{R}^2$ which does not pass through any of the points $\pm(\xi_0,\eta_0+2m\pi)$, $m\in\mathbf{Z}$. Let $n_m^\pm$ be the winding number of C with respect to $\pm(\xi_0,\eta_0+2m\pi)$. Then we have*

$$\begin{aligned}2\pi\sum_m &\left[n_m^+u(\xi_0,\eta_0+2m\pi)+n_m^-u(-\xi_0,-\eta_0-2m\pi)\right]\\ &=\int_C(u\partial_2v-v\partial_2u)\,d\xi+(v\partial_1u-u\partial_1v)\,d\eta,\end{aligned}$$

where ∂_1,∂_2 denote partial derivatives with respect to ξ, η, respectively.

In Theorem 5.1 we choose

$$u(\xi,\eta)=u_1(\xi)u_2(\eta),$$

where

$$u_1(\xi)=\mathrm{Ke}_n(\xi,h),\quad u_2(\eta)=\mathrm{ce}_n(\eta,q).$$

Let $\xi_0>0$ and $\eta_0\in\mathbf{R}$. We take the curve C to be the positively oriented boundary of the rectangle $\xi_1\le\xi\le\xi_2$, $\eta_0-\pi\le\eta\le\eta_0+\pi$, where $|\xi_1|<\xi_0<\xi_2$. Consider the line integral $\int_C$ in Theorem 5.1. Since u_2 has period 2π, the line integrals along the horizontal segments of C cancel each other. When $\xi_2\to+\infty$ the asymptotic behavior of $u_1(\xi)$ shows that the integral along the right-hand vertical segment of C tends to 0 as $\xi_2\to+\infty$. Therefore, setting

$$f(\xi)=\int_{-\pi}^{\pi}v(\xi,\eta,\xi_0,\eta_0)u_2(\eta)\,d\eta=\int_{\eta_0-\pi}^{\eta_0+\pi}v(\xi,\eta,\xi_0,\eta_0)u_2(\eta)\,d\eta$$

for $|\xi| < \xi_0$, one obtains

$$2\pi u_1(\xi_0)u_2(\eta_0) = u_1(\xi)f'(\xi) - u_1'(\xi)f(\xi). \tag{56}$$

We now argue as in section 2. By differentiating (56) with respect to ξ, we find that $f(\xi)$ satisfies the modified Mathieu equation (46). It is easy to see that $f(\xi)$ is an even function, so $f(\xi) = cu_3(\xi)$, where $u_3(\xi) = \mathrm{Ie}_n(\xi, h)$ and c is a constant. Then (56) implies that

$$2\pi u_1(\xi_0)u_2(\eta_0) = cW[u_1, u_3].$$

Since $W[u_1, u_3] = 1$,

$$\frac{1}{\pi}\int_{-\pi}^{\pi} v(\xi, \eta, \xi_0, \eta_0)u_2(\eta)\, d\eta = 2u_1(\xi_0)u_2(\eta_0)u_3(\xi) \quad \text{if } |\xi| < \xi_0. \tag{57}$$

By the same reasoning, we see that (57) is also true when $u_1(\xi) = \mathrm{Ko}_n(\xi, h)$, $u_2(\eta) = \mathrm{se}_n(\eta, q)$, $u_3(\xi) = \mathrm{Io}_n(\xi, h)$.

Expanding the function $\eta \mapsto v(\xi, \eta, \xi_0, \eta_0)$ according to Theorem 4.1, the following result is obtained.

Theorem 5.2. *Let $\xi, \eta, \xi_0, \eta_0 \in \mathbf{R}$ such that $|\xi| < \xi_0$, and let $k > 0$, $c > 0, q = -\frac{1}{4}c^2k^2$, $h = \frac{1}{2}ck$. Then*

$$\begin{aligned} K_0(kr) = {} & 2\sum_{n=0}^{\infty} \mathrm{Ie}_n(\xi, h)\mathrm{ce}_n(\eta, q)\mathrm{Ke}_n(\xi_0, h)\mathrm{ce}_n(\eta_0, q) \\ & + 2\sum_{n=1}^{\infty} \mathrm{Io}_n(\xi, h)\mathrm{se}_n(\eta, q)\mathrm{Ko}_n(\xi_0, h)\mathrm{se}_n(\eta_0, q), \end{aligned}$$

where r is given by (49).

Theorem 5.2 agrees with Meixner & Schäfke (1954) [14, section 2.66] although our notation and proof are different.

Inserting this result in (7), we obtain our final result.

Theorem 5.3. *Let $\mathbf{x}$, $\mathbf{x}_0$ be points on $\mathbf{R}^3$ with elliptic cylinder coordinates (ξ, η, z) and (ξ_0, η_0, z_0), respectively. If $\xi_{\lessgtr} := {\min \atop \max}\{\xi, \xi_0\}$ then*

$$\begin{aligned} \frac{1}{\|\mathbf{x} - \mathbf{x}_0\|} = {} & \frac{4}{\pi}\sum_{n=0}^{\infty}\int_0^{\infty} \mathrm{Ie}_n(\xi_<, h)\mathrm{ce}_n(\eta, q)\mathrm{Ke}_n(\xi_>, h)\mathrm{ce}_n(\eta_0, q)\cos k(z - z_0)\, dk \\ & + \frac{4}{\pi}\sum_{n=1}^{\infty}\int_0^{\infty} \mathrm{Io}_n(\xi_<, h)\mathrm{se}_n(\eta, q)\mathrm{Ko}_n(\xi_>, h)\mathrm{se}_n(\eta_0, q)\cos k(z - z_0)\, dk, \end{aligned}$$

where $q = -\frac{1}{4}c^2k^2$, $h = \frac{1}{2}ck$, and $\mathbf{x} \neq \mathbf{x}_0$.

Appendix A. Integrals for the (modified) parabolic cylinder harmonics expansion of $J_0(kr)$

The following formulas are valid for $\Im\lambda < 0$:

$$I_1 := \int_0^{\infty} J_0\left(\tfrac{1}{4}\xi^2\right) u_3(\lambda, \xi)\, d\xi \qquad = \frac{1}{2}\sqrt{\pi}(1 - i)\frac{G_1}{G_2^2}, \tag{A.1}$$

$$I_2 := \int_0^\infty \xi^{-1} J_1\left(\tfrac{1}{4}\xi^2\right) u_3(\lambda,\xi)\, d\xi = -\sqrt{\pi}\left(\lambda \frac{1}{G_2} + 2i\frac{G_2}{G_1^2}\right), \tag{A.2}$$

where

$$G_1 = G_1(\lambda) = \Gamma\left(\tfrac{1}{4} + \tfrac{i}{2}\lambda\right),$$
$$G_2 = G_2(\lambda) = \Gamma\left(\tfrac{3}{4} + \tfrac{i}{2}\lambda\right).$$

We believe that these integrals may be known but do not have a reference. We will derive (A.2). In (27) we use the integral representation [17, (13.4.4)]

$$\Gamma(a)U(a,b,z) = \int_0^\infty e^{-zt} t^{a-1}(1+t)^{b-a-1}\, dt, \quad \Re z, \Re a > 0.$$

Substituting $4s = \xi^2$ and changing the order of integration, one obtains

$$I_2 = \frac{1}{2G_1}\int_0^\infty t^{-\frac{3}{4}+\frac{i}{2}\lambda}(1+t)^{-\frac{3}{4}-\frac{i}{2}\lambda}\int_0^\infty s^{-1}J_1(s)e^{-is(2t+1)}\, ds\, dt. \tag{A.3}$$

From [22, page 405], we have, for $t > 0$,

$$\int_0^\infty s^{-1}J_1(s)\cos(s(2t+1))\, ds = 0,$$
$$\int_0^\infty s^{-1}J_1(s)\sin(s(2t+1))\, ds = t + (t+1) - 2\sqrt{t}\sqrt{t+1}.$$

Substituting these formulas in (A.3), we can evaluate I_2 using three times the formula for the beta function [17, (5.12.3)]

$$B(z,w) = \frac{\Gamma(z)\Gamma(w)}{\Gamma(z+w)} = \int_0^\infty t^{z-1}(1+t)^{-z-w}\, dt, \quad \Re z, \Re w > 0.$$

This gives (A.2).

The proof of (A.1) is similar, but in (27) one should first use [17, (13.2.40)]

$$U(a,b,z) = z^{1-b}U(1+a-b, 2-b, z).$$

The formulas (A.1), (A.2) remain valid for real λ. By separating real and imaginary parts, we obtain for $\lambda \in \mathbf{R}$,

$$\int_0^\infty J_0(\tfrac{1}{4}\xi^2)u_1(\lambda,\xi)\, d\xi = \frac{\Re(G_1\overline{G_2}) + \Im(G_1\overline{G_2})}{|G_2|^2}, \tag{A.4}$$

$$\int_0^\infty J_0(\tfrac{1}{4}\xi^2)u_2(\lambda,\xi)\, d\xi = \tfrac{1}{2}\left|\frac{G_1}{G_2}\right|^2, \tag{A.5}$$

$$\int_0^\infty \xi^{-1}J_1(\tfrac{1}{4}\xi^2)u_1(\lambda,\xi)\, d\xi = 2\left|\frac{G_2}{G_1}\right|^2 - \lambda, \tag{A.6}$$

$$\int_0^\infty \xi^{-1}J_1(\tfrac{1}{4}\xi^2)u_2(\lambda,\xi)\, d\xi = 2\frac{\Re(G_1\overline{G_2}) + \Im(G_1\overline{G_2})}{|G_1|^2},. \tag{A.7}$$

We may use

$$\Re(G_1\overline{G_2}) + \Im(G_1\overline{G_2}) = \frac{\pi\sqrt{2}e^{-\frac{1}{2}\pi\lambda}}{\cosh(\pi\lambda)}.$$

Formulas (A.4), (A.7) give us the integrals (35), (36) noting that we integrate even functions in (35), (36).

Acknowledgements

Part of this work was conducted while H. S. Cohl was a National Research Council Research Postdoctoral Associate in the Information Technology Laboratory at the National Institute of Standards and Technology, Gaithersburg, Maryland, U.S.A.

References

[1] F. V. Atkinson. *Discrete and continuous boundary problems.* Mathematics in Science and Engineering, Vol. 8. Academic Press, New York, 1964.

[2] H. Buchholz. *Die konfluente hypergeometrische Funktion mit besonderer Berücksichtigung ihrer Anwendungen.* Ergebnisse der angewandten Mathematik. Bd. 2. Springer-Verlag, Berlin, 1953.

[3] T. M. Cherry. Expansions in terms of parabolic cylinder functions. *Proceedings of the Edinburgh Mathematical Society. Series II*, 8:50–65, 1948.

[4] E. A. Coddington and N. Levinson. *Theory of ordinary differential equations.* McGraw-Hill Book Company, Inc., New York-Toronto-London, 1955.

[5] H. S. Cohl, J. E. Tohline, A. R. P. Rau, and H. M. Srivastava. Developments in determining the gravitational potential using toroidal functions. *Astronomische Nachrichten*, 321(5/6):363–372, 2000.

[6] C. G. Darwin. On Weber's function. *The Quarterly Journal of Mechanics and Applied Mathematics*, 2:311–320, 1949.

[7] A. Erdélyi, W. Magnus, F. Oberhettinger, and F. G. Tricomi. *Higher transcendental functions. Vol. II.* Robert E. Krieger Publishing Co. Inc., Melbourne, Fla., 1981.

[8] P. R. Garabedian. *Partial differential equations.* Chelsea Publishing Co., New York, second edition, 1986.

[9] E. Heine. *Handbuch der Kugelfunctionen, Theorie und Anwendungen (volume 2).* Druck und Verlag von G. Reimer, Berlin, 1881.

[10] E. W. Hobson. *The theory of spherical and ellipsoidal harmonics.* Chelsea Publishing Company, New York, 1955.

[11] N. N. Lebedev. *Special functions and their applications.* Dover Publications Inc., New York, 1972. Revised edition, translated from the Russian and edited by Richard A. Silverman, Unabridged and corrected republication.

[12] T. M. MacRobert. *Spherical Harmonics. An Elementary Treatise on Harmonic Functions with Applications.* Methuen & Co. Ltd., London, second edition, 1947.

[13] W. Magnus. Zur Theorie des zylindrisch-parabolischen Spiegels. *Zeitschrift für Physik. C. Particles and Fields*, 118:343–356, 1941.

[14] J. Meixner and F. W. Schäfke. *Mathieusche Funktionen und Sphäroidfunktionen mit Anwendungen auf physikalische und technische Probleme.* Die Grundlehren der mathematischen Wissenschaften in Einzeldarstellungen mit besonderer Berücksichtigung der Anwendungsgebiete, Band LXXI. Springer-Verlag, Berlin, 1954.

[15] J. C. P. Miller, editor. *Tables of Weber parabolic cylinder functions, giving solutions of the differential equation* $d^2y/dx^2+(\frac{1}{4}x^2-a)y=0$. Her Majesty's Stationery Office, London (British Information Services, 30 Rockefeller Plaza, New York 20, N. Y.), 1955. Computed by Scientific Computing Service Limited, Mathematical introduction by J. C. P. Miller.

[16] P. M. Morse and H. Feshbach. *Methods of theoretical physics. 2 volumes.* McGraw-Hill Book Co., Inc., New York, 1953.

[17] F. W. J. Olver, D. W. Lozier, R. F. Boisvert, and C. W. Clark, editors. *NIST handbook of mathematical functions.* Cambridge University Press, Cambridge, 2010.

[18] F. W. Schäfke. *Einführung in die Theorie der speziellen Funktionen der mathematischen Physik.* Die Grundlehren der mathematischen Wissenschaften, Bd. 118. Springer-Verlag, Berlin, 1963.

[19] H. Volkmer. Integralrelationen mit variablen Grenzen für spezielle Funktionen der mathematischen Physik. *Journal für die Reine und Angewandte Mathematik*, 319:118–132, 1980.

[20] H. Volkmer. Integralgleichungen für periodische Lösungen Hillscher differentialgleichungen. *Analysis. International Journal of Analysis and its Application*, 3(1-4):189–203, 1983.

[21] H. Volkmer. Integral representations for products of Lamé functions by use of fundamental solutions. *SIAM Journal on Mathematical Analysis*, 15(3):559–569, 1984.

[22] G. N. Watson. *A treatise on the theory of Bessel functions*. Cambridge Mathematical Library. Cambridge University Press, Cambridge, second edition, 1944.

[23] C. P. Wells and R. D. Spence. The parabolic cylinder functions. *Journal of Mathematics and Physics*, 24:51–64, 1945.